

Hierarchical Modelling for Spatiotemporal Data

Sudipto Banerjee¹ and Andrew O. Finley²

¹ Biostatistics, School of Public Health, University of Minnesota, Minneapolis, Minnesota, U.S.A.

² Department of Forestry & Department of Geography, Michigan State University, Lansing Michigan, U.S.A.

October 15, 2012



1

Specification:

- Again point-referenced vs. areal unit data
- Continuous time vs. discretized time

⇒ association in space, association in time

For point-referenced data, t continuous, Gaussian

$$Y(\mathbf{s}, t) = \mu(\mathbf{s}, t) + w(\mathbf{s}, t) + \epsilon(\mathbf{s}, t)$$

non-Gaussian data, $g(EY(\mathbf{s}, t) = \mu(\mathbf{s}, t) + w(\mathbf{s}, t))$

Don't treat time as a third coordinate (\mathbf{s}, t)

$$Cov(Y(\mathbf{s}, t), Y(\mathbf{s}', t')) = C(\mathbf{s} - \mathbf{s}', t - t')$$



2

- Separable form:

$$C(\mathbf{s} - \mathbf{s}', t - t') = \sigma^2 \rho_1(\mathbf{s} - \mathbf{s}'; \phi_1) \rho_2(t - t'; \phi_2)$$



3

- Separable form:

$$C(\mathbf{s} - \mathbf{s}', t - t') = \sigma^2 \rho_1(\mathbf{s} - \mathbf{s}'; \phi_1) \rho_2(t - t'; \phi_2)$$

- Nonseparable form:

- Sum of independent separable processes
- Mixing of separable covariance functions
- Spectral domain approaches



3

- Time discretized, $Y_t(\mathbf{s}), t = 1, 2, \dots, T$



4

- Time discretized, $Y_t(\mathbf{s}), t = 1, 2, \dots, T$
- Type of data: time series or cross-sectional



4

- Time discretized, $Y_t(\mathbf{s}), t = 1, 2, \dots, T$
- Type of data: time series or cross-sectional
- For time series data, exploratory analysis:

- Time discretized, $Y_t(\mathbf{s}), t = 1, 2, \dots, T$
- Type of data: time series or cross-sectional
- For time series data, exploratory analysis:
 - Arrange into an $n \times T$ matrix Y with entries $Y_t(\mathbf{s}_i)$

- Time discretized, $Y_t(\mathbf{s}), t = 1, 2, \dots, T$
- Type of data: time series or cross-sectional
- For time series data, exploratory analysis:
 - Arrange into an $n \times T$ matrix Y with entries $Y_t(\mathbf{s}_i)$
 - Center by row averages of Y yields Y_{rows}

- Time discretized, $Y_t(\mathbf{s}), t = 1, 2, \dots, T$
- Type of data: time series or cross-sectional
- For time series data, exploratory analysis:
 - Arrange into an $n \times T$ matrix Y with entries $Y_t(\mathbf{s}_i)$
 - Center by row averages of Y yields Y_{rows}
 - Center by column averages of Y yields Y_{cols}

- Time discretized, $Y_t(\mathbf{s}), t = 1, 2, \dots, T$
- Type of data: time series or cross-sectional
- For time series data, exploratory analysis:
 - Arrange into an $n \times T$ matrix Y with entries $Y_t(\mathbf{s}_i)$
 - Center by row averages of Y yields Y_{rows}
 - Center by column averages of Y yields Y_{cols}
 - sample spatial covariance matrix: $\frac{1}{T} Y_{rows} Y_{rows}^T$

- Time discretized, $Y_t(\mathbf{s}), t = 1, 2, \dots, T$
- Type of data: time series or cross-sectional
- For time series data, exploratory analysis:
 - Arrange into an $n \times T$ matrix Y with entries $Y_t(\mathbf{s}_i)$
 - Center by row averages of Y yields Y_{rows}
 - Center by column averages of Y yields Y_{cols}
 - sample spatial covariance matrix: $\frac{1}{T} Y_{rows} Y_{rows}^T$
 - sample autocorrelation matrix: $\frac{1}{n} Y_{cols}^T Y_{cols}$

- Time discretized, $Y_t(\mathbf{s}), t = 1, 2, \dots, T$
- Type of data: time series or cross-sectional
- For time series data, exploratory analysis:
 - Arrange into an $n \times T$ matrix Y with entries $Y_t(\mathbf{s}_i)$
 - Center by row averages of Y yields Y_{rows}
 - Center by column averages of Y yields Y_{cols}
 - sample spatial covariance matrix: $\frac{1}{T} Y_{rows} Y_{rows}^T$
 - sample autocorrelation matrix: $\frac{1}{n} Y_{cols}^T Y_{cols}$
 - E , residuals matrix after a regression fitting, Empirical orthogonal functions (EOF)

- **Modeling:** $Y_t(\mathbf{s}) = \mu_t(\mathbf{s}) + w_t(\mathbf{s}) + \epsilon_t(\mathbf{s})$, or perhaps $g(E(Y_t(\mathbf{s}))) = \mu_t(\mathbf{s}) + w_t(\mathbf{s})$

- **Modeling:** $Y_t(\mathbf{s}) = \mu_t(\mathbf{s}) + w_t(\mathbf{s}) + \epsilon_t(\mathbf{s})$, or perhaps $g(E(Y_t(\mathbf{s}))) = \mu_t(\mathbf{s}) + w_t(\mathbf{s})$
- For $\epsilon_t(\mathbf{s})$, i.i.d. $N(0, \tau_t^2)$

- **Modeling:** $Y_t(\mathbf{s}) = \mu_t(\mathbf{s}) + w_t(\mathbf{s}) + \epsilon_t(\mathbf{s})$, or perhaps $g(E(Y_t(\mathbf{s}))) = \mu_t(\mathbf{s}) + w_t(\mathbf{s})$
- For $\epsilon_t(\mathbf{s})$, i.i.d. $N(0, \tau_t^2)$
- For $w_t(\mathbf{s})$
 - $w_t(\mathbf{s}) = \alpha_t + w(\mathbf{s})$
 - $w_t(\mathbf{s})$ independent for each t
 - $w_t(\mathbf{s}) = w_{t-1}(\mathbf{s}) + \eta_t(\mathbf{s})$, independent spatial process innovations

Dynamic spatiotemporal models

Measurement Equation

$$Y(\mathbf{s}, t) = \mu(\mathbf{s}, t) + \epsilon(\mathbf{s}, t); \quad \epsilon(\mathbf{s}, t) \stackrel{ind}{\sim} N(0, \sigma_\epsilon^2).$$

$$\mu(\mathbf{s}, t) = \mathbf{x}(\mathbf{s}, t)' \tilde{\beta}(\mathbf{s}, t).$$

$$\tilde{\beta}(\mathbf{s}, t) = \beta_t + \beta(\mathbf{s}, t)$$

Transition Equation

$$\beta_t = \beta_{t-1} + \eta_t, \quad \eta_t \stackrel{ind}{\sim} N_p(\mathbf{0}, \Sigma_\eta)$$

$$\beta(\mathbf{s}, t) = \beta(\mathbf{s}, t-1) + \mathbf{w}(\mathbf{s}, t).$$